

**INFLUENCE OF PRIMARY COSMIC RADIATION MASS COMPOSITION
ON THE ESTIMATION OF EAS ENERGY**S. P. KNURENKO*, A. A. IVANOV, V. A. KOLOSOV, Z. E. PETROV,
I. YE. SLEPTSOV and G. G. STRUCHKOV*Yu. G. Shafer Institute of Cosmophysical Research and Aeronomy Lenin Avenue 31, Yakutsk 677891, Russia***s.p.knurenko@ikfia.ysn.ru*

Received (Day Month Year)

Revised (Day Month Year)

Energy portion E_{em}/E_0 transferred to EAS electron–photon component at $E_0 = 10^{15} \pm 10^{19}$ eV was estimated by using of Čerenkov radiation data and charged particles data obtained at the Yakutsk array. The results are compared with the models for different energy dissipation into EAS electron–photon component and with calculations for different primary nuclei. In the energy intervals $10^{15} \pm 10^{16}$ eV and $10^{18} \pm 10^{19}$ eV E_{em}/E_0 is equal to (77 ± 2) and $(88 \pm 2)\%$ respectively, that doesn't contradict with mixed composition of primary particles for the first energy interval and the proton composition for the second one.

Keywords: Cosmic Rays; Extensive Air Showers; Mass Composition.

1. Introduction

Energy characteristics of a shower such as the energy lose by particles on ionization of medium and total energy transferred to electron–photon EAS component are always considered as important for choice of the interaction mechanism of a cosmic ray primary particle with nuclei of atoms of the air. And these characteristics can be used for estimation of primary particle energy without a definite model. The energy can be estimated only in two cases (without using of any notions of the interaction model): when the total flux of Čerenkov light at the sea level has been measured; the longitudinal EAS development, a more exactly, the total number of particles in the maximum of shower development has been measured. At present time, only at two EAS arrays such measurements are carried out: at the Yakutsk array (Russia) where the Čerenkov EAS radiation is measured and at the HiRes array (USA) where measurements of the ionization nitrogen luminosity are provided.

In the present work, a problem of determination of the energy portion transferred to electromagnetic EAS cascade is studied. The wide energy interval from 10^{15} to 10^{19} eV is considered. In the method of determination of E_{em} the real conditions of the atmosphere, the longitudinal development of a shower (X_{max}) and the mass composition of primary particles are taken into account.

2 S. P. Knurenko et al.

2. Energy Transferred to the Electromagnetic EAS Component

Figure 1 presents the Yakutsk EAS array experimental data together with calculations by the model with decelerated and moderate dissipation of energy into electromagnetic EAS component: quasiscaling models (solid line) and QGSJET (dashed line)¹. From Fig. 1




Fig. 1. A portion of the energy transferred to the electromagnetic EAS component by Čerenkov light data at the Yakutsk array.

one can see both the agreement of experimental data with calculations by QGSJET model (proton) in the region $E_0 \geq 3 \times 10^{18}$ eV, and disagreement at $E_0 \leq 3 \times 10^{18}$ eV. The scaling model gives a noticeably greater value of E_{em}/E_0 in relation to the experimental data that is doubtlessly also connected with the break of scaling function in the region of ultra-high energies.

The experimental data in Fig. 1 is well approximated by the expression of a form:

$$E_{em}/E_0 = (0.964 \pm 0.011) - (0.079 \pm 0.005) \times E_0^{-(0.147 \pm 0.008)}, \quad (1)$$

where E_0 is taken in units of eV.

The relation (1) is primarily important for the comparison of estimations of E_0 obtained at the Yakutsk and Fly's Eye arrays. The new estimation of E_{cal}/E_0 parameter for the Fly's Eye array is given in Ref. 2:

$$E_{cal}/E_0 = (0.959 \pm 0.003) - (0.082 \pm 0.003) \times E_0^{-(0.147 \pm 0.006)} \quad (2)$$

From (1) and (2) a good coincidence of both results, calculation and empirical estimation is well seen. The calculations in Ref. 1 have been carried out by using of QGSJET model in the case of primary proton and iron nucleus (see Table 1). In the case of the pri-

Table 1. Yakutsk data and calculations for p/Fe

n/n	E_0 , eV	experiment E_{em}/E_0	QGSJET model E_{em}/E_0	SIBYLL model E_{em}/E_0
1	10^{17}	0.824 ± 0.015	0.876/0.806	—
2	10^{18}	0.872 ± 0.008	0.901/0.850	0.915/0.865
3	10^{19}	0.911 ± 0.010	0.914/0.879	0.934/0.897

mary proton a good agreement between both calculations is observed. The comparison of experimental data with calculations for the proton and iron nucleus indicates the fact that the mass compositions of cosmic radiation particles in the energy region of $10^{17} - 10^{19}$ eV and above 3×10^{18} eV must differ. At $E_0 \geq 3 \times 10^{18}$ eV the mass composition is most likely close to the proton one. Therefore, when estimating the EAS energy at the Fly's Eye array it is more reasonably to use the formula (2). At the Yakutsk array the formula (1) obtained empirically is used. A systematic difference between (1) and (2) does not exceed 10% that speaks for a coincidence of estimations of E_0 at the Yakutsk and Fly's Eye arrays.

Acknowledgements

This work has been financially supported by RFBR, grant #02–02–16380, grant #03–02–17160 and grant INTAS #03–51–5112.

References

1. S. P. Knurenko *et al.*, in *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City* **1**, 372 (1999).
2. C. Song, Z. Cao, B. R. Dawson *et al.*, *Astropart. Phys.* **14**, 7 (2000).